arXiv:1412.1689v1 [math.GM] 3 Dec 2014

On an alternative proof of Fermat's Last Theorem

Mike Winkler

December 3, 2014

Abstract

Conjecture: The quadratic Diophantine equation system

$$\begin{aligned} q^2 &= a^2\alpha - b^2\beta - c^2\gamma, \\ pq &= (ad)^2\alpha - (be)^2\beta - (cf)^2\gamma, \\ p^2 &= (ad^2)^2\alpha - (be^2)^2\beta - (cf^2)^2\gamma, \end{aligned}$$

has no nontrivial integer solution for $d \neq e \neq f \neq 0$, if there is $\alpha = \beta = \gamma = 1$, or $|\alpha| \neq |\beta| \neq |\gamma| \neq 0$ with $\alpha|a,\ \beta|b,\ \gamma|c$ and $|\alpha| \neq a,\ |\beta| \neq b,\ |\gamma| \neq c$.

Fermat's Last Theorem states that the Diophantine equation $x^n + y^n = z^n$ has no nontrivial integer solution (x, y, z) for all $n > 2$. A proof was published in 1995 by Andrew Wiles[4]. It is shown that a proof of the above conjecture proves Fermat's Last Theorem for all $n > 5$.

A possible alternative proof

Lemma: Let

$$r = x - y, \qquad s = y + z, \qquad t = z + x,$$

and

$$u = x + y + z, \qquad v = y - z - x, \qquad w = x - y - z,$$

there is with

$$\begin{aligned} \mathcal{A} &= r^2(u^4 - 1)(xy)^{n-2} - s^2(v^4 - 1)(yz)^{n-2} - t^2(w^4 - 1)(zx)^{n-2}, \\ \mathcal{B} &= 2(ru)^2(xy)^{n-2} - 2(sv)^2(yz)^{n-2} - 2(tw)^2(zx)^{n-2}, \\ \mathcal{C} &= r^2(u^4 + 1)(xy)^{n-2} - s^2(v^4 + 1)(yz)^{n-2} - t^2(w^4 + 1)(zx)^{n-2}, \end{aligned}$$

the identity

$$(8rst)^2(xyz)^{n-2}(x^n + y^n - z^n) = \mathcal{A}^2 + \mathcal{B}^2 - \mathcal{C}^2. \tag{1}$$

Proof: Let set $r = x - y,\ s = y + z,\ t = z + x$ and $u = x + y + z,\ v = y - z - x,\ w = x - y - z$ into (1). After multiplying out and summarizing we get an equation with two identical sides. □

Theorem: The Diophantine equation $x^n + y^n = z^n$ has no nontrivial integer solution (x, y, z) for all $n > 5$.

Proof: Let (x, y, z) be a nontrivial integer solution for a specific $n > 2$ of

$$x^n + y^n = z^n. \tag{2}$$

It is sufficient only to look at the solutions (x, y, z) with $gcd(x, y, z) = 1$. Therewith instantly follows the coprimeness of x, y, z. Of the numbers x, y, z is then exactly one even. With a solution (x, y, z) for (2) the left side of (1) comes equal to zero, so there is

$$\mathcal{A}^2 + \mathcal{B}^2 = \mathcal{C}^2 \tag{3}$$

All integer solutions for (3) can be represented by integers p, q with $p > q > 0$ as follows:

$$\mathcal{A} = p^2 - q^2, \qquad \mathcal{B} = 2pq, \qquad \mathcal{C} = p^2 + q^2.$$

According to the Lemma, there is

$$p^2 - q^2 = r^2(u^4 - 1)(xy)^{n-2} - s^2(v^4 - 1)(yz)^{n-2} - t^2(w^4 - 1)(zx)^{n-2}, \tag{4}$$

$$2pq = 2(ru)^2(xy)^{n-2} - 2(sv)^2(yz)^{n-2} - 2(tw)^2(zx)^{n-2}, \tag{5}$$

$$p^2 + q^2 = r^2(u^4 + 1)(xy)^{n-2} - s^2(v^4 + 1)(yz)^{n-2} - t^2(w^4 + 1)(zx)^{n-2}. \tag{6}$$

From subtraction of (4) and (6) it follows

$$q^2 = r^2(xy)^{n-2} - s^2(yz)^{n-2} - t^2(zx)^{n-2}. \tag{7}$$

From (5) is followed by division by 2

$$pq = (ru)^2(xy)^{n-2} - (sv)^2(yz)^{n-2} - (tw)^2(zx)^{n-2}. \tag{8}$$

From addition of (4) and (6) it follows

$$p^2 = (ru^2)^2(xy)^{n-2} - (sv^2)^2(yz)^{n-2} - (tw^2)^2(zx)^{n-2}. \tag{9}$$

Now we consider separately the cases for even and odd $n > 2$.

Let $n = 2k + 1$ with $k \in \mathbb{N}$, we get with (7), (8), (9) the equation system

$$\begin{aligned} q^2 &= \left(r(xy)^{k-1}\right)^2 xy - \left(s(yz)^{k-1}\right)^2 yz - \left(t(zx)^{k-1}\right)^2 zx, \\ pq &= \left(ru(xy)^{k-1}\right)^2 xy - \left(sv(yz)^{k-1}\right)^2 yz - \left(tw(zx)^{k-1}\right)^2 zx, \\ p^2 &= \left(ru^2(xy)^{k-1}\right)^2 xy - \left(sv^2(yz)^{k-1}\right)^2 yz - \left(tw^2(zx)^{k-1}\right)^2 zx. \end{aligned} \tag{10}$$

Let $a = r(xy)^{k-1}$, $b = s(yz)^{k-1}$, $c = t(zx)^{k-1}$ and $d = u$, $e = v$, $f = w$ and $\alpha = xy$, $\beta = yz$, $\gamma = zx$, we get with (10) the equation system of the conjecture. For the non-existence of integer solutions there must $d \neq e \neq f \neq 0$ and $|\alpha| \neq |\beta| \neq |\gamma| \neq 0$ and $\alpha|a$, $\beta|b$, $\gamma|c$ and $|\alpha| \neq a$, $|\beta| \neq b$, $|\gamma| \neq c$.

According to the Lemma, there is $r = x - y$, $s = y + z$, $t = z + x$ and $u = x + y + z$, $v = y - z - x$, $w = x - y - z$. Therefore we verify the conditions of the conjecture.

For a nontrivial integer solution (x, y, z) of (2) we can assume that $|x| \neq |y| \neq |z| \neq 0$.

Let $|x| \neq |y| \neq |z| \neq 0$, there is $x + y + z \neq y - z - x \neq x - y - z \neq 0$. So, there is $u \neq v \neq w \neq 0$ and $d \neq e \neq f \neq 0$.

Let $|x| \neq |y| \neq |z| \neq 0$, there is $|xy| \neq |yz| \neq |zx| \neq 0$ and so $|\alpha| \neq |\beta| \neq |\gamma| \neq 0$.

For all $k > 1$ there is $xy|r(xy)^{k-1}, yz|s(yz)^{k-1}, zx|t(zx)^{k-1}$, and so $\alpha|a$, $\beta|b$, $\gamma|c$.

Let $x, y, z \in \mathbb{Z}$ with $gcd(x, y, z) = 1$ and $|x| \neq |y| \neq |z| \neq 0$, there is $x - y \neq y + z \neq z + x \neq 0$, and so $r \neq s \neq t \neq 0$. For all $k > 2$ there is $|xy| \neq r(xy)^{k-1}, |yz| \neq s(yz)^{k-1}, |zx| \neq t(zx)^{k-1}$, and so $|\alpha| \neq a$, $|\beta| \neq b$, $|\gamma| \neq c$.

Because for all $k > 2$ the conditions of the conjecture are complied, it follows that (10) has no nontrivial integer solution (p, q, x, y, z) for all $k > 2$.

Let $n = 2k$ with $k \in \mathbb{N}, k > 1$, we get with (7), (8), (9) the equation system

$$\begin{aligned} q^2 &= \left(r(xy)^{k-1}\right)^2 - \left(s(yz)^{k-1}\right)^2 - \left(t(zx)^{k-1}\right)^2, \\ pq &= \left(ru(xy)^{k-1}\right)^2 - \left(sv(yz)^{k-1}\right)^2 - \left(tw(zx)^{k-1}\right)^2, \\ p^2 &= \left(ru^2(xy)^{k-1}\right)^2 - \left(sv^2(yz)^{k-1}\right)^2 - \left(tw^2(zx)^{k-1}\right)^2. \end{aligned} \tag{11}$$

For the equation system (11) the same conditions apply as for (10), except that $\alpha = \beta = \gamma = 1$. Therefore even here for all $k > 1$ the conditions of the conjecture are complied. From this it follows directly that (11) has no nontrivial integer solution (p, q, x, y, z) for all $k > 1$.

Under the assumption of the validity of the conjecture, equation (2) has no nontrivial integer solution (x, y, z) for all $n > 5$, what proves the Theorem. □

References

1. Fillipi, Andreas: Post in the online forum Matroids Matheplanet.
(http://www.matheplanet.de/matheplanet/nuke/html/viewtopic.php?topic=192975)

2. Scheid, Harald: Zahlentheorie. 3.Auflage. Spektrum Akademischer Verlag, Heidelberg u.a. 2003, ISBN 3-8274-1356-6.

3. Wikipedia: Fermat's Last Theorem.
(http://en.wikipedia.org/wiki/Fermat's_Last_Theorem)

4. Wiles, Andrew: Modular Elliptic Curves and Fermat's last theorem. Annals of Mathematics 142 (1995), S. 443551.
(http://math.stanford.edu/∼lekheng/flt/wiles.pdf)

5. Mike Winkler, Über die Lösbarkeit der diophantischen Gleichung $x^n + y^n = z^n$, April 2014.
(http://mikewinkler.co.nf/fermat_n>2_2014.pdf)

This article is a translation of my German article[5].

Mike Winkler, Ernst-Abbe-Weg 4, 45657 Recklinghausen, Germany

mike.winkler(at)gmx.de